

ON THE RESIDUAL SOLVABILITY OF GENERALIZED FREE PRODUCTS OF FINITELY GENERATED NILPOTENT GROUPS

D.KAHROBAEI

Dedicated to Gilbert Baumslag on the occasion of his 70th Birthday

ABSTRACT. In this paper we study the residual solvability of the generalized free product of finitely generated nilpotent groups. We show that these kinds of structures are often residually solvable.

1. INTRODUCTION AND MOTIVATION

The notion of residual properties was first introduced by Philip Hall in 1954 [9]. Let X be a class of groups. G is residually- X if, for every non-identity element g in G , there is an epimorph of G in X such that the element corresponding to g is not the identity.

In this paper we focus on residual solvability. The notion of residual solvability of groups was pioneered particularly by Gilbert Baumslag in his celebrated paper [4] where he proved that positive one-relator groups are residually solvable. The author in [7] studied this notion in general. In [6] the author studied residual solvability of residually solvable groups.

In 1963 Gilbert Baumslag studied residual finiteness of generalized free products of finitely generated torsion-free nilpotent groups [1]. He showed first that if both factors are non-abelian then this amalgamation is not residually-finite and he found conditions that ensure G is residually finite. He actually proved in general that this kind of structure is free-by-residually-finite, or meta-residually-finite. A few years later, in 1968, one of his students, Joan Landman Dyer, continued his work [8] on residual finiteness of these structures. She showed in particular that

Date: February 8, 2008.

2000 Mathematics Subject Classification. Primary 20F19.

Key words and phrases. residual solvability, generalized free products, nilpotent groups, abelianization, generalized central products, poly-residual solvability.

The research of this author has been supported by CUNY research foundation at the City College of the City University of New York and New York Group Theory Cooperative.

if the factors are not finitely generated nilpotent groups, these kinds of groups are not free-by-residually-finite (by taking isomorphic factors of class three), but rather are residually-finite-by-free-by-residually-finite.

It is interesting to mention that the generalized free product of two finitely generated nilpotent groups (or two finitely generated free groups) with a finitely generated subgroup amalgamated has a solvable word problem (see page 150 [5]). In this work we consider the generalized free product of finitely generated nilpotent groups, and discuss how close such groups are to being residual solvable. We show how conditions on the amalgamating subgroup affects residual solvability. Note that the free products of finitely generated nilpotent groups are residually solvable.

Effect of abelianization on amalgamated products of nilpotent groups

We give a complete explanation of abelianization of generalized free products. In particular, we see how the abelianization of such groups with nilpotent factors is not trivial, using the Frattini theorem, which states that the commutator subgroup of a finitely generated nilpotent group is a non-generator set. As a result, we conclude that generalized free products of two finitely generated nilpotent groups are not perfect. Note that the condition of being finitely generated is required in these results since we use the Frattini theorem.

Theorem 1. *The generalized free product of two finitely generated nilpotent groups amalgamated by a proper subgroup of them is not perfect.*

This property of finitely generated nilpotent groups plays a critical role in our approach to our question. For example finitely generated polycyclic factors that satisfy the maximal condition, but fail to satisfy the condition of the Frattini theorem, turn out to be perfect under certain conditions.

Cyclic amalgamated subgroup and residual solvability

We first consider the case when the amalgamating subgroup is cyclic. It turns out that choosing an appropriate solvable filtration of each factor so that the generator of amalgamating subgroup does not lie in the n -th term of upper central series enables us to prove that it is residually solvable-by-solvable.

Theorem 2. *The generalized free product of two finitely generated nilpotent groups amalgamated by a cyclic subgroup is residually solvable.*

The amalgamated subgroup is central in all factors

We next define the generalized central product of an arbitrary number of groups, and show that each factor injects into such a product. We then look at the case where the amalgamating subgroup is central in all factors, and we show using the generalized central products that these structures turn out to be free-by-nilpotent. We note that a nilpotent extension of a free group is not residually nilpotent but rather is residually solvable.

Theorem 3. *The generalized free product of an arbitrary number of finitely generated nilpotent groups of bounded class, amalgamating a central subgroup in each of the factors, is an extension of a free group by a nilpotent group. Furthermore, such groups are residually solvable.*

The case where one of the factors is abelian

We then consider the case where just one of the factors is abelian. It turns out that such groups with an abelian factor are free-by-nilpotent.

Theorem 4. *The generalized free product of a finitely generated torsion-free abelian and a nilpotent group is residually solvable-by-abelian-by-finite abelian. Furthermore such groups are residually solvable.*

The case when the amalgamated subgroup is of finite index in at least one of the factors

Theorem 5. *The generalized free product of two finitely generated torsion-free nilpotent groups amalgamating a proper subgroup which is of finite index in at least one of the factors, is an extension of a free group by a torsion-free nilpotent group. Furthermore, such groups are residually solvable.*

Note that here the condition of being torsion-free is necessary for the factors, because of Mal'cev's Fundamental Theorem, as we see later.

Doubles of nilpotent groups and residual solvability, arbitrary number of factors

By a double we mean the amalgamated product of two groups where the factors are isomorphic and the amalgamated subgroups are identified under the same isomorphism. One can generalize the definition of doubles to arbitrary number of factors.

Theorem 6. *Let $\{A_i | i \in I\}$ be an arbitrary indexed family of isomorphic torsion-free nilpotent groups, such that $\bigcap_{i \in I} A_i = C$, and let G be the generalized free product of A_i s amalgamated by C :*

$$G = \left\{ \prod_{i \in I}^* A_i; C \right\}.$$

Then G is an extension of a free by nilpotent group. Furthermore, G is residually solvable.

Generalized free products of nilpotent groups sometimes may fail to be residually solvable.

Gilbert Baumslag gave a counter example in [3] that shows that not every subgroup of the generalized free product of two finitely generated torsion-free nilpotent groups is indicable. We use this example to prove the following proposition:

Proposition 7. *There exist two finitely generated torsion-free non-abelian free nilpotent groups A and B such that the amalgamated product of them with abelian amalgamating subgroup $C_A = C_B$,*

$$G = \{A * B; C_A = C_B\},$$

is not residually solvable and not poly-residually solvable.

This implies that an abelian amalgamating subgroup is not sufficient for residual solvability.

Poly-residual solvability of the generalized free product of nilpotent groups

As Proposition 7 suggests, we need to impose some conditions on the amalgamating subgroup to ensure that the generalized free product of finitely generated nilpotent group be poly-residually solvable; by a poly-residually solvable group we mean a group that has at least one poly-residually solvable series (see Section 12 for precise definition). Here is the theorem that gives us these conditions:

Theorem 8. *The generalized free product of two finitely generated nilpotent groups, A and B amalgamating subgroups of them, C_A , C_B respectively is poly-residually solvable if the following condition holds: the solvable filtrations of A and B which are defined on the upper central series of A and B respectively, are solvable with compatible filtrations, i.e. $\xi_i A \cap C_A \stackrel{\phi_i}{=} \xi_i B \cap C_B$.*

Acknowledgment. I thank my Ph.D. supervisor G.Baumslag and also K.J.Falconer and P.de la Harpe for helpful comments.

2. BACKGROUND AND PRELIMINARY RESULTS

In this section we recall some definitions and facts and prove some lemmas to be used later.

Recall G is called an extension of A by Q if there exist A and Q and a short exact sequence $1 \rightarrow A \rightarrow G \rightarrow Q \rightarrow 1$. We say G is meta- X ,

where X is a property (or class), if G is an extension of A by Q where A and Q have property (or class) X .

2.1. Subgroups of amalgamated products. We will use a theorem of Hanna Neumann [13] extensively in this paper. With regard to abstract groups, Hanna Neumann showed in the 1950s that, in general, subgroups of amalgamated products are no longer amalgamated products, but generalized free products, indeed she proved the following: let K be a subgroup of $G = \{A * B; C\}$, then K is an HNN-extension of a tree product in which the vertex groups are conjugates of subgroups of either A or B and the edge groups are conjugates of subgroups of C . The associated subgroups involved in the HNN-extension are also conjugates of subgroups of C . If K misses the factors A and B (i.e. $K \cap A = \{1\} = K \cap B$), then K is free; and if K misses the amalgamated subgroup C (i.e. $K \cap C = \{1\}$), then $K = \prod_{i \in I}^* X_i * F$, where the X_i are conjugates of subgroups of A and B and F is free (see [5] for more information).

Let us mention that later a description was given by the Bass-Serre theory [16], with groups acting on graphs to give geometric intuition: the fundamental group of a graph of groups generalizes both amalgamated products, HNN-extensions and tree products.

3. SOME RESULTS ON THE STRUCTURE OF THE ABELIANIZATION

The following lemma formulates the abelianization of the amalgamated products of two groups. (We leave it as an exercise to the reader to check this.)

Lemma 9. *Let G be the amalgamated product of two groups A and B amalgamating C ,*

$$G = \{A * B; C\}.$$

Then the abelianization of G_{ab} of G takes the following form:

$$G_{ab} = (A_{ab} \times B_{ab}) / gp(\bar{c}\alpha\bar{c}^{-1}\beta | c \in C),$$

where α is the monomorphism from C into A and β is the monomorphism from C into B .

Lemma 10. *Let G be the amalgamated product of A and B identifying C_A with C_B ,*

$$G = \{A * B; C_A = C_B\}.$$

Then the abelianization G_{ab} of G maps onto

$$D = A_{ab} / gp(\bar{c}_a | c_a \in C_A) \times B_{ab} / gp(\bar{c}_b^{-1} | c_b \in C_B).$$

Proof. By Lemma 9, the abelianization of G can be expressed as:

$$G_{ab} = (A_{ab} \times B_{ab})/gp(\bar{c}_a \bar{c}_b^{-1}).$$

Put

$$\begin{aligned} N &= gp(\bar{c}_a \bar{c}_b^{-1}) \\ N_1 &= gp(\bar{c}_a) \\ N_2 &= gp(\bar{c}_b^{-1}). \end{aligned}$$

Let θ be the map from G_{ab} into D ,

$$\begin{aligned} \theta : G_{ab} &\rightarrow D \\ \theta : (A_{ab} \times B_{ab})/N &\rightarrow A_{ab}/N_1 \times B_{ab}/N_2 \end{aligned}$$

defined by:

$$\bar{a}N \mapsto \bar{a}N_1 \text{ and } \bar{b}N \mapsto \bar{b}N_2.$$

A typical element w in N takes the form $w = \bar{c}_a \bar{c}_b^{-1}$, so

$$w = \bar{c}_a \bar{c}_b^{-1} \mapsto (\bar{c}_a N_1, \bar{c}_b^{-1} N_2) = (N_1, N_2).$$

and by Von Dyck's theorem θ is an epimorphism as requested. $\square$

The following is an alternative proof of Lemma 10, using presentations.

Proof. Let A and B have presentations:

$$A = \langle X; R \rangle \text{ and } B = \langle Y; S \rangle.$$

So the abelianizations of A and B have presentations:

$$\begin{aligned} A_{ab} &= \langle X; R \cup \{[x_i, x_j] \mid x_i, x_j \in X\} \rangle \\ B_{ab} &= \langle Y; S \cup \{[y_i, y_j] \mid y_i, y_j \in Y\} \rangle. \end{aligned}$$

The direct product of A_{ab} and B_{ab} has the following presentation since, by definition of the direct product, all elements of A_{ab} commute with all elements of B_{ab} :

$$\begin{aligned} A_{ab} \times B_{ab} &= \langle X \cup Y; \quad R \cup S \cup \{[x_i, x_j] \mid x_i, x_j \in X\} \\ &\quad \cup \{[y_i, y_j] \mid y_i, y_j \in Y\} \\ &\quad \cup \{[x, y] \mid x \in X, y \in Y\} \rangle. \end{aligned}$$

With α the monomorphism from C into A , and β the monomorphism from C into B , put

$$\begin{aligned} N &= gp(c\alpha \ c\beta^{-1} \mid c \in C) \\ N_1 &= gp(c\alpha \mid c \in C) \\ N_2 &= gp(c\beta^{-1} \mid c \in C). \end{aligned}$$

By Lemma 9, G_{ab} , the abelianization of G , has the following presentation:

$$\begin{aligned} G_{ab} = (A_{ab} \times B_{ab})/N = \langle X \cup Y; R \cup S \cup & \{[x_i, x_j] | x_i, x_j \in X\} \\ & \cup \{[y_i, y_j] | y_i, y_j \in Y\} \\ & \cup \{[x, y] | x \in X, y \in Y\} \\ & \cup \{c\alpha c\beta^{-1} | c \in C\} \rangle. \end{aligned}$$

Then the presentation of each factor of D is

$$\begin{aligned} A_{ab}/N_1 &= \langle X; R \cup \{[x_i, x_j] | x_i, x_j \in X\} \cup \{c\alpha | c \in C\} \rangle \\ B_{ab}/N_2 &= \langle Y; S \cup \{[y_i, y_j] | y_i, y_j \in Y\} \cup \{c\beta^{-1} | c \in C\} \rangle. \end{aligned}$$

Their direct product D has the following presentation:

$$\begin{aligned} D = A_{ab}/N_1 \times B_{ab}/N_2 = & \langle X \cup Y; R \cup S \cup \{[x_i, x_j] | x_i, x_j \in X\} \\ & \cup \{[y_i, y_j] | y_i, y_j \in Y\} \\ & \cup \{c\alpha | c \in C\} \\ & \cup \{c\beta^{-1} | c \in C\} \\ & \cup \{[x(c\alpha), y(c\beta^{-1})] | x \in X, y \in Y, c \in C\} \rangle. \end{aligned}$$

Now define θ to be a map from G_{ab} into D , by sending $x_i \mapsto x_i$ and $y_i \mapsto y_i$. Then $N \mapsto 1$, so by Von Dyck's Theorem, θ defines a homomorphism from G_{ab} onto D . $\square$

4. EFFECTS OF THE ORDER OF THE ABELIANIZATION OF AMALGAMATED PRODUCTS OF NILPOTENT GROUPS

In order to study the effect of the order of the abelianization of the amalgamated products of nilpotent groups, we first study the effect of indicability on the order of finitely generated nilpotent groups. Recall that a group A is termed indicable if there exists a homomorphism of A onto the infinite cyclic group. A finitely generated group G is indicable if and only if G_{ab} is infinite. Higman [10] proves that every finitely generated torsion-free nilpotent group is indicable and is therefore infinite. We conclude that the abelianization of every finitely generated torsion-free nilpotent group is also infinite. Note that the abelianization of an infinite finitely generated nilpotent group A is again infinite, since there is a canonical homomorphism from A onto $A/\tau A$, where τA is the torsion group of A . Since $A/\tau A$ is a finitely generated torsion-free nilpotent group, $A/\tau A$ is infinite. Hence A is infinite.

To prove the following proposition we use a theorem of G. Baumslag [3] that states that the amalgamated product of two finitely generated torsion-free nilpotent groups is indicable.

Proposition 11. *The abelianization of the amalgamated product of two finitely generated nilpotent groups is infinite only if one of the factors is infinite.*

Proof. First let A and B be two finitely generated torsion-free nilpotent groups, let C be a proper subgroup of both of them, and G be the amalgamated product of them, amalgamating C ,

$$G = \{A * B; C\}.$$

G is indicable by the theorem of G.Baumslag. Therefore G_{ab} , the abelianization of G , is infinite (see [5] for more information). Now let just one of the factors be torsion-free. There exists an epimorphism from G onto a finitely generated torsion-free nilpotent group,

$$G^* = \{A/\tau A * B/\tau B; C/\tau C\},$$

the amalgamated product of finitely generated torsion-free nilpotent groups. Therefore G^*_{ab} is infinite, using the first case, implying that G_{ab} is infinite. $\square$

5. PROOF OF THEOREM 1

In this section we first recall the Frattini Subgroup Theorem. Recall that the Frattini subgroup of G , ΦG , is the intersection of all maximal subgroups of G . In particular, ΦG is characteristic. We also recall that $g \in G$ is called a non-generator of G , if whenever $G = gp(X, g)$ then $G = gp(X)$. A theorem of Frattini states that ΦG is the set of all non-generators in G . In particular if G is a finitely generated nilpotent group, then $\delta_2 G \leq \Phi G$, where $\delta_2 G = [G, G]$ (see [14] for more information).

Proof. Let A and B be two finitely generated nilpotent groups, and let C_A and C_B be proper subgroups of A and B respectively. We want to show that the generalized free product $G = \{A * B; C_A = C_B\}$ is not perfect, that is that the abelianization G_{ab} of G is not trivial.

By Lemma 10, G_{ab} maps onto

$$D = A_{ab}/gp(\bar{c}_a | c_a \in C_A) \times B_{ab}/gp(\bar{c}_b^{-1} | c_b \in C_B).$$

where $\bar{c}_a$ and $\bar{c}_b$ are the images of $c_a \in C_A$ and $c_b \in C_B$ in A_{ab} and B_{ab} respectively.

We claim that under the conditions of the Theorem D is not trivial. In order to prove the claim, we will show that

$$A_{ab}/gp(\bar{c}_a) \neq \{1\} \quad \text{and} \quad B_{ab}/gp(\bar{c}_b^{-1}) \neq \{1\}.$$

Since C_A is a proper subgroup of A , the Frattini Subgroup Theorem implies that $gp(C_A, \delta_2 A)$ is a proper subgroup of A , where $\delta_2 A = [A, A]$.

Since C_A is a proper subgroup of A , $gp(C_A, \delta_2 A)$ is also a proper subgroup of A , so

$$A/gp(C_A, \delta_2 A) \neq \{1\}.$$

By using the third isomorphism theorem we have:

$$\frac{A_{ab}}{gp(\bar{c}_a)} \simeq \frac{\frac{A}{\delta_2 A}}{\frac{gp(C_A, \delta_2 A)}{\delta_2 A}} \simeq \frac{A}{gp(C_A, \delta_2 A)} \neq \{1\}.$$

Similarly

$$B_{ab}/gp(\bar{c}_b^{-1}) \neq \{1\}.$$

This proves our claim that, $D \neq \{1\}$ and so $G_{ab} \neq \{1\}$. Hence G is not perfect. $\square$

6. PROOF OF THEOREM 2

Proof. Let A and B be two finitely generated nilpotent groups. Let a be a non-identity element of A , and b be a non-identity element of B . We can find $m \geq 1$ and $n \geq 1$ such that $1 \neq a \in \xi_{m+1}A \setminus \xi_m A$, and $1 \neq b \in \xi_{n+1}B \setminus \xi_n B$, (where $\xi_i A$ is the i -th term of the upper central series of A , and $\xi_j B$ is the j -th term of the upper central series of B). Let $G = \{A * B; a = b\}$ be the generalized free product of A and B amalgamating a with b . Let D be the central product of $A/\xi_m A$ and $B/\xi_n B$ amalgamating $a\xi_m A$ with $b\xi_n B$,

$$D = \{A/\xi_m A \times B/\xi_n B; a\xi_m A = b\xi_n B\}.$$

Note that

$$C\xi_n B/\xi_n B \simeq C/C \cap \xi_n B$$

and

$$C\xi_m A/\xi_m A \simeq C/C \cap \xi_m A$$

are cyclic groups. This confirms the fact that

$$a\xi_m A = b\xi_n B.$$

Map G into D and let K be the kernel of this map. Then observe that $K \cap C = \{1\}$, where $C = gp(a) = gp(b)$. Therefore K is a free product of conjugates of subgroups of A and B , and a free group by the Hanna Neumann Theorem, see Section 2.1. So K is residually solvable. Hence G is an extension of a residually solvable group by a solvable group. $\square$

7. THE AMALGAMATED SUBGROUP IS CENTRAL IN ALL FACTORS

7.1. Generalized central products and some related results.

Here we introduce a general definition of the generalized central product of an arbitrary number of factors. Let us mention the work of D.Robinson in [14] for finite number of factors. We adopt the notation used for generalized free products [5]:

Definition 12.

Suppose that

$$\{A_i = \langle X_i; R_i \rangle | i \in I\}$$

is an indexed family of presentations

$$A_i = \langle X_i; R_i \rangle$$

of the groups A_i , and suppose C is another group equipped with monomorphisms

$\phi_i : C \rightarrow A_i$ and $C \leq \xi A_i$ (for all $i \in I$, where ξA_i is the center of A_i).

We term the group A defined by the presentation

$$A = \langle \cup X_i; \cup R_i \cup \{c\phi_i c^{-1}\phi_j | c \in C, i, j \in I\} \cup \{[x_i, x_j] = 1, i, j \in I\} \rangle,$$

where we assume that the X_i are disjoint, the generalized central product of A_i amalgamating the central subgroup C .

$$A = \prod_{i \in I}^{\times} \{A_i; C\}.$$

If $C = 1$, then A is termed the direct product of the A_i .

According to Von Dyck's theorem, there are canonical homomorphisms μ_i of each A_i to A . We will prove that the μ_i are monomorphisms, and, if we identify A_i with $A_i\mu_i$, then

$$c\phi_i = c\phi_j \text{ for all } c \in C, i, j \in I.$$

Thus, we can identify C with any of its images $C\phi_i$ which are already identified with $C\phi_j\mu_i$. Then $A_i \cap A_j = C$ ($i, j \in I, i \neq j$) and $A = gp(\cup_{i \in I} A_i)$. A can also be written as

$$A = (\prod_{i \in I}^{\times} A_i) / gp(c\phi_i c^{-1}\phi_j | i, j \in I, c \in C).$$

Lemma 13. $\mu_i : A_i \rightarrow (\prod_{i \in I}^{\times} A_i) / gp(c\phi_i c^{-1}\phi_j | i, j \in I, c \in C)$ is a monomorphism for all $i \in I$.

Proof. Put $S = gp(c\phi_i c^{-1}\phi_j | i, j \in I, c \in C)$, where $C \xrightarrow{\phi_i} A_i$. To show that μ_i is a monomorphism we must show that $\ker \mu_i = 1$. Let $a \in A_i$ and $a \in \ker \mu_i$, so that $a\mu_i = 1$. We want to show that $a = 1$. A generic element in S has the following form:

$$s = (c_1\phi_{i_1}c_1^{-1}\phi_{j_1}) \cdots (c_n\phi_{i_n}c_n^{-1}\phi_{j_n}).$$

Since $a\mu_i = 1$, then $a \in S$, and

$$a = (c_1\phi_{i_1}c_1^{-1}\phi_{j_1}) \cdots (c_n\phi_{i_n}c_n^{-1}\phi_{j_n}).$$

Let us consider two cases: the case where none of i_r and j_r are equal to i , and the case where some of them are equal to i .

Case 1: Since none of the subscripts are equal to i , this implies that

$$a \in gp(A_k | k \neq i).$$

But a is also an element in A_i . Therefore

$$a \in A_i \cap gp(A_k | k \neq i) = 1$$

(by a property of the direct product), so $a = 1$.

Case 2: Now suppose that some of the indexes are equal to i , say $i_l = i$.

Note that for $c \in C$, $c\phi_i$ is central in A_i . Thus we have:

$$a = \left(\prod_{i_l} c_l\phi_{i_l}c_l^{-1}\phi_{j_l} \right) \left(\prod_{k_m \neq i} c_1\phi_{k_1}c_1^{-1}\phi_{k_1} \cdots c_n\phi_{k_n}c_n^{-1}\phi_{k_n} \right).$$

By a similar argument,

$$\left(\prod_{k_m \neq i} c_1\phi_{k_1}c_1^{-1}\phi_{k_1} \cdots c_n\phi_{k_n}c_n^{-1}\phi_{k_n} \right) = 1.$$

Now

$$\prod_{i_l} c_l\phi_{i_l} = \left(\prod c_l \right) \phi_i = c\phi_i.$$

Hence

$$a = (c\phi_i) \left(\prod_{j_l \neq i} c_l^{-1}\phi_{j_l} \right).$$

We have

$$ac^{-1}\phi_i = \left(\prod_{j_l \neq i} c_l^{-1}\phi_{j_l} \right) \in A_i \cap gp(A_k | k \neq i) = 1,$$

so $a = c\phi_i$. This implies that $a = 1$, since

$$a = c\phi_i = \left(\prod_{i_l} c_l\phi_{i_l}c_l^{-1}\phi_{j_l} \right).$$

The right hand side has an even number of factors, so this is possible only if it is equal to 1, as required. $\square$

7.2. Proof of Theorem 3. Before proving the Theorem, we bring together some of the facts and related lemmas that we will use in the proof. One of these facts is that a direct product of finitely many nilpotent groups is nilpotent. However we note that a direct product of infinitely many nilpotent groups is not necessarily nilpotent. (see D.Segal's book [15] for more information)

Lemma 14. *The generalized central product of finitely many nilpotent groups is nilpotent.*

Proof. The generalized central product of nilpotent groups is the quotient of the direct product of finitely many nilpotent groups, (which is again nilpotent, [15] page 6) and another nilpotent group. The quotient of nilpotent groups is nilpotent. Therefore the generalized central product of finitely many nilpotent groups is nilpotent. $\square$

We note that the generalized central product of finitely many solvable groups is solvable. For the case of abelian factors, the generalized central product of an arbitrary number of abelian groups is abelian. Now we prove Theorem 3:

Proof. Suppose that $\{A_i | i \in I\}$ is an indexed family of finitely generated nilpotent groups and let G be the generalized free product of the A_i , with amalgamating subgroup C :

$$G = \left\{ \prod_{i \in I}^* A_i; C \right\}.$$

Let S be the generalized central product of A_i (see Definition 12):

$$S = \prod_{i \in I}^{\times} A_i / gp(c\phi_i c^{-1}\phi_j | i, j \in I, c \in C).$$

S contains a canonical homomorphism of A_i for all $i \in I$. Since these homomorphisms coincide on C , they can be extended to a homomorphism μ from G into S . Let K be the kernel of μ . Since S is nilpotent, it follows that G/K is nilpotent. By Lemma 13, μ is one-to-one restricted to each factor, i.e.

$$K \cap A_i = 1 \text{ for all } i \in I.$$

So, by the theorem of Hanna Neumann mentioned in Section 2.1, K is free. Hence G is a nilpotent extension of a free group, so is also residually solvable. $\square$

The following proposition is for the case of two finitely generated solvable factors.

Proposition 15. *The generalized free product of two finitely generated solvable groups amalgamated by central subgroups, is a solvable extension of a residually solvable group. Furthermore such groups are residually solvable.*

Proof. Let N be the normal closure of B in G , i.e. $N = gp_G(B)$. Since A/C is solvable and $G/N \simeq A/C$ then G/N is a solvable group. Since N is an iterated, infinite untwisted double of copies of B , which therefore can be mapped onto B with free kernel, N is residually solvable. So G is a solvable extension of a residually solvable group. $\square$

For the case of abelian factors, we have the following corollary:

Corollary 16. *The generalized free product an of arbitrary number of abelian groups is an extension of a free group by an abelian group. Further, such groups are residually solvable.*

8. PROOF OF THEOREM 4

Here we prove Theorem 4. Note that this is a special instance of the case where the amalgamating subgroup is central in only one of the factors.

Proof. Let A and B be finitely generated groups, with A torsion-free abelian and B torsion-free nilpotent, and let G be their amalgamated product of them with amalgamating subgroup C ,

$$G = \{A * B; C_A = C_B\}.$$

Then C is a direct factor of a subgroup A_1 of A of finite index, i.e.

$$A_1 = C \times H \quad \text{where } [A : A_1] = n < \infty.$$

There is an epimorphism from G onto $A/A_1 = \bigcup_{i=1}^n a_i A_1$, where the a_i are a distinct set of coset representatives of A_1 in A . The kernel of this map is $K = gp_G(B, A_1) = \{\prod_{i=1}^n {}^* B^{a_i} * A_1; C\}$. Now let D be the normal closure of finitely many copies of B in K :

$$D = gp_K(\bigcup_{i=1}^n B^{a_i}) = \{\prod_{i=1}^n {}^* B^{a_i}; C\}.$$

Since $[a_i, C] = 1$, then D is the generalized free product of a finite number of doubles, and hence D is residually solvable. So K is an extension of a residually solvable group by an abelian group. Therefore G is residually solvable-by-abelian-by-finite abelian. Therefore G is residually solvable. $\square$

8.1. An example where one of the factors is abelian. Here we construct an example to illustrate the above theorem.

Example 17. Let $A = gp(a, b, c)$ be a free abelian group, and $B = gp(x, y, z)$ be a free nilpotent group of class 2. $C_A = gp(a^2, b)$ and $C_B = gp(y, z)$ are free abelian groups of rank 2. If we form

$$G = \{A * B; a^2 = x, b = z\},$$

then G is residually solvable. Note that C_B is normal but not central in B .

Proof. The presentations of A , B , C_A and C_B are as follows:

$$A = \langle a, b, c; [a, b], [b, c], [a, c] \rangle \text{ free of rank 3,}$$

$$C_A = gp(a^2, b) = \langle a^2, b; [a^2, b] \rangle \text{ free abelian of rank 2,}$$

$$B = \langle x, y, z; [x, y] = z, [x, z], [y, z] \rangle \text{ free nilpotent of class 2 and of rank 3,}$$

$$C_B = gp(y, z) = \langle y, z; [y, z] \rangle \text{ free abelian of rank 2.}$$

Define an isomorphism ϕ which maps

$$a^2 \mapsto y \text{ and } b \mapsto z.$$

Form the generalized free product of A and B identifying C_A with C_B :

$$G = \{A * B; C_A = C_B\}.$$

C_B is normal in B but is not central in B , and observe that $\xi B \cap C_B \neq \{1\}$.

However, C_A is central and therefore normal in A . On the other hand the quotient group B/C_B is an infinite cyclic group generated by xC_B ,

$$B/C_B = gp(xC_B).$$

Let K be the normal closure of A in G ,

$$K = gp_G(A) = gp_B(A) = gp(A^{x^i} | i \in \mathbb{Z}).$$

G/K is an infinite cyclic group,

$$G/K = gp(xK).$$

Note that

$$a^{x^i} = b^i a^2 \text{ and } b^{x^i} = b \text{ for all } i \in \mathbb{Z}.$$

Hence

$$A \cap A^{x^i} = C_A \quad (\forall i \in \mathbb{Z})$$

so defining

$$A_i := gp(A, A^{x^i}) = \{A * A^{x^i}; C_A\}$$

we have

$$\forall w_i \in A_i \exists w_j \in A_j \text{ s.t. } w_i = w_j^{x^{(i-j)}}$$

Thus

$$K = \langle A_i, x^i (i \in \mathbb{Z}); H_i = H_j^{x^{(i-j)}} (i, j \in \mathbb{Z}) \rangle.$$

Then K is an HNN-extension, where the base groups are the union of A_i ($i \in \mathbb{Z}$), the stable letters are $x_i = x^i$ ($i \in \mathbb{Z}$), and the associated subgroups are H_i , where H_i is a subgroup of A_i (for $i \in \mathbb{Z}$). In particular by using the subgroup theorem, we can express K as the generalized free product of conjugates of free abelian groups.

$$K = \left\{ \prod_{i \in \mathbb{Z}}^* A^{x^i}; C_A \right\}.$$

Hence K is an extension of a free group by an abelian group. Consider the following short exact sequence:

$$1 \rightarrow K \rightarrow G \rightarrow B/C_B \simeq \mathbb{Z} \rightarrow 1.$$

G/K is an infinite cyclic group, so therefore G is an extension of a residually solvable group by an infinite cyclic group, and thus G is residually solvable. $\square$

9. PROOF OF THEOREM 5

We first give some background we need to prove Theorem 5.

9.1. The Mal'cev completion and the fundamental theorem of torsion-free nilpotent groups. Here we recall some definitions and theorems that we use to prove Theorem 5. A group is called complete if, for every element a of G and natural number n , the equation $x^n = a$ has at least one solution in G , or, in other words, every root of every element of G belongs to G . There is a theorem that states that a finitely generated nilpotent group is complete if and only if it contains no proper subgroup of finite index; hence every group is contained in a complete group. Let $\mathcal{D}$ denote the set of the class of groups in which extraction of roots is always uniquely possible. Let us term a minimal torsion-free nilpotent $\mathcal{D}$ -group, $m(G)$, containing a given torsion-free nilpotent group G , a Mal'cev completion of G .

Let G be a finitely generated torsion-free nilpotent group of class c , let g be an element of G and let n be a positive integer. Then G can be embedded in a finitely generated torsion-free nilpotent group of class c in which g has an n -th root. Therefore every finitely generated torsion-free nilpotent group can be embedded in a nilpotent $\mathcal{D}$ -group.

We recall the Mal'cev Theorem: let A be a torsion-free nilpotent group, then A can be embedded in a torsion-free nilpotent $\mathcal{D}$ -group. As a corollary we have: let G, H be torsion-free nilpotent groups, let ϕ be a homomorphism of G into H and let $m(G)$ and $m(H)$ be any Mal'cev completions of G and H ; then ϕ can be extended uniquely to a homomorphism $m(\phi)$ of $m(G)$ onto $m(H)$. Also if $m(G)$ and $m'(G)$ are Mal'cev completions of G then they are isomorphic. We now state Mal'cev's fundamental theorem of torsion-free nilpotent groups. If G_1^* and G_2^* are two completions of G , then there exists an isomorphism between them that extends the identity automorphism of G , and this isomorphism is unique (for more details see [2], [11]).

9.2. Proof of Theorem 5. Note that here the condition of being torsion-free is necessary, because of Mal'cev's Fundamental Theorem.

Proof. Let A and B be two finitely generated torsion-free nilpotent groups, and let C be a proper subgroup of A and B , such that the index of C in, say, A is finite. We want to show that the generalized free product of A and B with C amalgamating,

$$G = \{A * B; C\},$$

is a finitely generated torsion-free nilpotent extension of a free group. Let $m(A)$ and $m(B)$ be Mal'cev completions of A and B respectively. Since the index of C in A is finite, then $m(A)$ is also a Mal'cev completion of C . There exists a monomorphism μ from C into B . Then μ can be extended uniquely to a homomorphism $m(\mu)$ of $m(A)$ onto $m(B)$. Now, there is a homomorphism from G into $m(B)$, which is consistent on C , and injective when restricted to A and B . Let K be the kernel of this homomorphism, then we have

$$K \cap A = \{1\} = K \cap B.$$

Therefore K is free by the result of Hanna Neumann mentioned in Section 2.1. Hence G is a torsion-free nilpotent extension of a free group. $\square$

10. PROOF OF THEOREM 6

We first need to prove the following lemma which will be used in proving Theorem 6.

Lemma 18. *If A is a group, C is a subgroup of A , ϕ is an isomorphic mapping of A onto a group B , and D is the amalgamated product of A and B amalgamating C with $C\phi$, that is*

$$D = \{A * B; C = C\phi\},$$

then there is a homomorphism, ψ , from D onto one of the factors, with kernel:

$$\ker \psi = gp(a(a\phi)^{-1} | a \in A).$$

Furthermore ψ injects into each factor.

Proof. Let α be the homomorphism from A onto itself, and β be the homomorphism from B onto the inverse of the isomorphic copy of A , i.e. $\beta = \phi^{-1}$. These homomorphisms can be extended to a homomorphism from D onto A , ([5] page 103, [12]). Since $\psi(Cab) = Ca(b\phi^{-1})$ it follows easily $K = \ker \psi = gp(a(a\phi^{-1}) | a \in A)$. By the way that α and β are defined, it follows that this homomorphism is one-to-one restricted to either A or B . $\square$

Now we are ready to prove Theorem 6.

Proof. For each $i \in I$ let $\phi : G \rightarrow A_i$ be an epimorphism, and let K be the kernel of ϕ . K is free, since ϕ restricted to each factor is injective, and

$$A_i \cap K = \{1\} \quad (\forall i \in I).$$

Therefore G is free-by-nilpotent. $\square$

11. PROOF OF PROPOSITION 7

Proof. Let $A = gp(a, x)$ be a free nilpotent group of class 2, and $B = gp(b, y)$ be a free nilpotent group of class 3. Then $C_A = gp(a, a^x)$ and $C_B = gp(b, [b, b^y])$ are free abelian groups of rank 2. If we form

$$\begin{aligned} G &= \{A * B; C_A = C_B\} \\ &= \{A * B; a = b, a^x = [b, b^y]\}, \end{aligned}$$

then

$$a = [a, a^y]^x = [a^x, a^{y^x}],$$

so a lies in every term of the derived series of G . Hence G is not residually solvable. Now put $N = gp_G(a)$ which is perfect. Since a poly-residually solvable groups can not have a perfect subgroup, G is not poly-residually solvable. $\square$

Note that this contrasts with the case where residual finiteness of the groups has been studied, where Baumslag proved that generalized free products of finitely generated torsion-free nilpotent groups in general are meta-residually solvable.

We have shown that if the amalgamating subgroup of the generalized free product of torsion-free nilpotent groups is cyclic (Proposition 2) or central in both factors (Theorem 15) then these groups are residually

solvable. One may conjecture that it would be the case when the amalgamating subgroup is abelian. However this is false by the example of Baumslag as mentioned above.

This can be easily shown using Baumslag's example of an amalgamated product, which can be modified to give an example where the factors are finite, but the product is not residually solvable.

12. PROOF OF THEOREM 8

Let us recall the definition of a poly-property. Let X be a property (or class) of groups. A finite normal series

$$1 = G_0 \leq G_1 \leq \cdots \leq G_l = G \quad (1)$$

of G is termed a poly- X series for G if

$$G_{i+1}/G_i \in X \text{ for } i = 0, \dots, l-1.$$

A group G is termed poly- X if it has at least one poly- X series. The length of the series (1) is l .

The proof of Theorem 8 uses an upper central filtration approach:

Proof. Let $G = \{A * B; C_A = C_B\}$ be the generalized free product of finitely generated nilpotent groups A and B . Assume that $\xi_i A$ and $\xi_i B$ be i -th terms of the upper central series of A and B respectively and $(\xi_i A), (\xi_i B)$ (for i is bounded by the maximum class of nilpotency of A and B) be solvable filtrations of A and B such that they are also compatible filtrations. We want to show that G has an invariant series

$$1 = G_0 \leq G_1 \leq \cdots \leq G_k = G \quad (k < \infty)$$

such that G_{i+1}/G_i is residually solvable (for $i = 0, \dots, k-1$). Map

$$G \rightarrow G/gp_G(C_A \cap \xi_1 A).$$

Note that, since $\xi G = \xi A \cap \xi B \cap C_A$, then

$$gp_G(C_A \cap \xi_1 A) < \xi_1 G.$$

$$\begin{aligned} G/gp_G(C_A \cap \xi_1 A) &\simeq \{A/gp_G(C_A \cap \xi_1 A) * B/gp_G(C_A \cap \xi_1 A); \\ &\quad C_A/gp_G(C_A \cap \xi_1 A) = C_B/gp_G(C_A \cap \xi_1 A)\} \\ G/gp_G(C_A \cap \xi_1 A) &\simeq \{A/C_A \cap \xi_1 A * B/C_B \cap \xi_1 B; \\ &\quad C_A/C_A \cap \xi_1 A = C_B/C_B \cap \xi_1 B\}. \end{aligned}$$

One can check that this is residually solvable.

$$\begin{aligned} G/gp_G(C_A \cap \xi_1 A)/gp_G(\xi_1 A/C_A \cap \xi_1 A, \xi_1 B/C_B \cap \xi_1 B) &\simeq \\ \{A/\xi_1 A * B/\xi_1 B; C_A \xi_1 A/\xi_1 A = C_B \xi_1 B/\xi_1 B\}. \end{aligned}$$

Inductively we can show that for i each quotient is residually solvable and this completes the proof of the theorem. $\square$

REFERENCES

1. G. Baumslag, *On the residual finiteness of generalised free products of nilpotent groups*, Trans. of Amer. Math. Soc. **106** (1963), 193–209.
2. ———, *Lecture notes on nilpotent groups*, Regional Conference Series in Mathematics, American Mathematical Society, Providence, R.I. **2** (1971), 1–73.
3. ———, *On generalised free product of torsion-free nilpotent groups*, Illinois J. Math. **16** (1971), 526–528.
4. ———, *Positive one-relator groups*, Trans. of Amer. Math. Soc. **156** (1971), 165–183.
5. ———, *Topics in combinatorial group theory*, Birkhauser-Verlag, 1993.
6. D.Kahrobaei, *Are doubles of residually solvable groups residually solvable?*, Preprint.
7. ———, *Residual solvability of generalized free products*, The City University of New York (GC) PhD Thesis, October 2004.
8. J. Landman Dyer, *On the residual finiteness of generalized free products*, Trans. Amer. Math. Soc. **133** (1968), 131–143.
9. P. Hall, *The splitting properties of relatively free groups*, Proc. Lond. Math. Soc. **4** (1954), 343–356.
10. G. Higman, *The units of group-rings*, Proc. Lond. Math. Soc. **46** (1940), 231–246.
11. A. G. Kurosh, *The theory of groups*, Chelsea Publishing Company, 1960.
12. B. H. Neumann, *An essay on free products of groups with amalgamations*, Philos. Trans. Roy. Soc. London. Ser. A. **246** (1954), 503–554.
13. H. Neumann, *Generalized free products with amalgamated subgroups*, Amer. J. Math. **71** (1949), 491–540.
14. D.J.S. Robinson, *A course in the theory of groups*, Springer-Verlag, New York, 1995.
15. D. Segal, *Polycyclic groups*, Cambridge University Press, 1983.
16. J.-P. Serre, *Trees*, Springer-Verlag, 1980.

MATHEMATICAL INSTITUTE, UNIVERSITY OF ST ANDREWS, NORTH HAUGH,
ST ANDREWS, FIFE, KY16 9SS, UK

E-mail address: delaram.kahrobaei@st-andrews.ac.uk

URL: <http://www-groups.mcs.st-and.ac.uk/~delaram/>